

Finnrick Analytics
Finnrick.com



Project 145165
Lab # 203663
Date Rec'd 4/7/2026
Report Issued

Certificate of Analysis

Compound Retatrutide

Label Claim 5 mg

Batch/Lot f2rnhkw



Analyte

Peptide Analysis

Chromatographic purity

Retatrutide

Mass Spectral Match

Microbiology

Endotoxins

FDA Critical Metals

Arsenic

Cadmium

Lead

Mercury

<u>Result</u>	<u>Units</u>	<u>LOQ</u>	<u>% of Label</u>	<u>Method</u>	<u>Date</u>	<u>CAS</u>
>99.9	%	0.5		HPLC-UV/MS	4/13/2026	
4.75	mg	0.5	95.0	HPLC-UV-MS	4/13/2026	2381089-83-2
ID Confirmed				HPLC-UV-MS	4/13/2026	
<0.5	EU	0.5		USP 86	4/14/2026	
ND	µg	0.50		USP 233	4/19/2026	7440-38-2
ND	µg	0.10		USP 233	4/19/2026	7440-43-9
0.55	µg	0.50		USP 233	4/19/2026	7439-92-1
ND	µg	0.10		USP 233	4/19/2026	7439-97-6

The data presented are from the analysis of the sample shown and meet Krause Analytical internal quality assurance criteria unless otherwise flagged.

Methods shown reference current Krause Analytical SOPs

ND - Not detected LOQ - limit of quantification

All values reported on a per vial basis unless otherwise noted

Copyright © 2026 Krause Analytical LLC. All rights reserved.

Respectfully submitted,

Mark C. Krause
Laboratory Director

8127 Mesa Drive Suite B-206 Austin, TX 78759

CHROMATOGRAM




Finnrick
Safer.

Openlynx Report
File:203663

Sample Report:

1: (Time: 0.30) Combine (13:19-(7:10+29:32))

1:MS ES+
4.3e+005



**krause analytical**

Methods Summary

Purity/Potency/Identification

USP/NF 621

- 2 mL of purified water is added to the lyophilized powder in the vial, and the contents mixed to dissolve the lyophilized powder.
- An aliquot is taken from the vial and diluted to contain approximately 500 mg/L of the peptide.
- The diluted sample is analyzed by HPLC-UV-MS.
- The mass spectrum obtained is compared to an authentic standard of the peptide for identification.
- The total area of all of the peaks in the chromatogram is calculated, and the area of the peak of the peptide is divided by the total area to obtain the chromatographic purity value, reported in percent.
- The area of the peptide is compared to the area of the peptide peak in the known standard to obtain a concentration in the solution. This concentration is used to calculate the total mass of peptide in the vial, which is compared to the stated mass (label claim) and reported as both total mass in the vial and as a percent of the label claim.

Endotoxins

USP/NF 85

- 2 mL of purified water is added to the lyophilized powder in the vial, and the contents mixed to dissolve the lyophilized powder.
- An aliquot is taken from the vial and diluted in endotoxin-free water.
- The diluted sample is analyzed for endotoxins using the LAL method.

Metals

USP/NF 233

- 2 mL of purified water is added to the lyophilized powder in the vial, and the contents mixed to dissolve the lyophilized powder.
- An aliquot is taken from the vial and diluted in deionized water.
- The diluted aliquot is analyzed against known standards by ICP-MS

Krause Analytical, LLC 8127 Mesa Drive Suite B-206 Austin, TX 78759